

**ANNE GRADY SERVICES RESPITE DEPARTMENT
PEDIATRIC STANDING ORDERS (Pg 1 of 2)**

NAME _____ DATE _____

These orders need to be signed by the child's Primary Care Physician, or Waterville Family Physicians. The orders may be initiated by the wing nurse in consultation with the supervisory nurse or by the supervisory nurse after careful assessment of the child staying here. When used, they must be brought forward to the active physician's orders and recorded for 72 hours. Orders contraindicated should have a line drawn through them. All orders are PRN. All oral medications should be given only per G-tube if NPO and receives only enteral feedings.

FOR FEVER OR DISCOMFORT:

TYLENOL 80MG/0.8 ML DROPS

6-11 lbs½ dppr/0.4 ml every 4 hours
12-17 lbs1 dppr/0.8 ml every 4 hours
18-23 lbs..... 1 ½ dppr/1.2 ml every 4 hours
24-35 lbs2 dppr/1.6 ml every 4 hours

TYLENOL 160 MG/5 ML ELIXIR

6-11 lbs..... ¼ tsp. every 4 hours
12-17 lbs. ½ tsp. every 4 hours
18-23 lbs... ¾ tsp. every 4 hours
24-35 lbs.....1 tsp. every 4 hours
36-47 lbs.....1 ½ tsp. every 4 hours
48-59 lbs..... 2 tsp. every 4 hours
60-71 lbs.....2 ½ tsp. every 4 hours
72-95 lbs3 tsp. every 4 hours
96 lbs & over.... 4 tsp. every 4 hours

IBUPROFEN INFANT DROPS 50 MG/1.25 ML

12-17 lbs.....1.25 ml every 6-8 hours
18-23 lbs.....1.875 ml every 6-8 hours

IBUPROFEN ORAL SUSPENSION 100 MG/5 ML

12-17 lbs... ½ tsp. every 6-8 hours
18-23 lbs..... ¾ tsp. every 6-8 hours
24-35 lbs 1 tsp. every 6-8 hours
36-47 lbs..... 1 ½ tsp. every 6-8 hours
48-59 lbs 2 tsp. every 6-8 hours
60-71 lbs..... 2 ½ tsp. every 6-8 hours
72 -95 lbs..... 3 tsp. every 6-8 hours

TYLENOL 80 MG CHEWABLE TABS

24-35 lbs2 tabs every 4 hours
36-47 lbs3 tabs every 4 hours
48-59 lbs4 tabs every 4 hours
60-71 lbs5 tabs every 4 hours
72-95 lbs.....6 tabs every 4 hours

TYLENOL 325 MG TABLETS

48-95 lbs 1 tab every 4 hours
96 lbs & over 2 tabs every 4 hours

IBUPROFEN JR STRENGTH 100 MG TABLETS

48-71 lbs.....2 tabs every 6-8 hours
72-95 lbs.....3 tabs every 6-8 hours

IBUPROFEN 200 MG TABLETS

48-71 lbs..... 1 tab every 6-8 hours
72 lbs & over..... 2 tabs every 6-8 hours

FOR BEGINNING SIGNS OF NASAL CONGESTION:

SUDAFED LIQUID 30 MG/5 ML

<2 years 4 mg/kg/day in divided doses every 6 hours (1 kg = 2.2 lbs)
2-6 years ½ tsp. every 4-6 hours
6-11 years 1 tsp. every 4-6 hours
12 years and older 2 tsp. every 4-6 hours

FOR COUGHS:

ROBITUSSIN DM LIQUID 100 MG/5 ML

<2 years 1-2mg/kg/day in divided doses every 6-8 hours (1kg = 2.2 lbs)
2-6 years ½ tsp. every 4 hours
6-11 years 1 tsp. every 4 hours
12 years and older 2 tsp. every 4 hours

PEDIATRIC STANDING ORDERS (Pg 2 of 2)

FOR CONSTIPATION:

GLYCERIN SUPPOSITORIES (After 3 days without BM)

1-5 years 1 suppository
6 years and older 2 suppositories

DULCOLAX SUPPOSITORIES (After 4 days without BM)

1-2 years ½ suppository
2 years and older 1 suppository

FLEETS ENEMA

1-12 years 1 pediatric enema
12 years and older 1 adult enema

FOR DIARRHEA:

KAOPECTATE LIQUID

Under 1 year 1 tsp. three times a day
1-6 years 2 tsp four times a day

IMMODIUM ORAL SOLUTION 1 MG/5 ML

6-8 years (48-59 lbs) 2 tsp. after 1st loose stool
1 tsp. after each loose stool
thereafter (MAX/day – 4 tsp)
9-11 years (60-98 lbs) 2 tsp. after 1st loose stool
1 tsp. after each loose stool
thereafter (MAX/day – 6 tsp)
12 years and older 4 tsp. after 1st loose stool
2 tsp. after each loose stool
thereafter (MAX/day – 8 tsp)

IMMODIUM 2MG TABLET

6-11 years 1 tab
12 years and older 2 tabs

FOR MINOR ABRASIONS:

1. Cleanse with Hydrogen Peroxide 5%
2. Apply triple antibiotic ointment twice a day
3. Notify the physician if necessary

FOR SKIN IRRITATION:

Apply A & D ointment to affected area twice a day

FOR BUG BITES/AREAS CAUSING ITCHING:

Apply Caladryl or Cortaid lotion topically to affected area 3 times a day

FOR NOSEBLEED:

Squeeze nose, tilt head forward. If persists, apply cold compress to nose.

Physician Signature X _____

Date _____

Guardian Signature X _____

Date _____